

Ridin' Herd

► by **Rick Rasby**, Extension beef specialist, University of Nebraska

Keep heifer development costs in check

Cow-calf producers have experienced some of the best times in recent history of the beef industry. Total calf inventory has been low, and demand for calves has been high, resulting in good calf prices.

Anyone who has been in this industry for any length of time understands the cattle industry is a cyclic industry, where cow numbers expand and contract and cattle prices fluctuate. Even in good times, it is critical to explore opportunities to reduce costs. In considering nutrition programs for the young beef female, I'm reminded of the statement the late Jim Wiltbank emphasized

in his talks: "If you are going to breed 'em, you have to feed 'em."

It has routinely been recommended that heifers be developed to two-thirds of their mature weight before the start of the breeding season, because puberty in beef females is a function of weight. In addition, in most cow herds, the lowest pregnancy rate occurs in females that are asked to conceive

their second calf, and underdeveloped heifers at first breeding may be ones that are in a continual catch-up mode and are challenged for their next pregnancy.

Lighter prebreeding weights

Because the replacement heifer enterprise does not generate income, or very little income from cull replacements, costs for this enterprise are borne by the cow-calf enterprise. Our data suggest replacement heifer costs from the time the replacement heifer is weaned to the beginning of her first calving season can add \$69-\$88 to the annual cow costs. The more economical the replacement heifer program, the greater the profit potential of the cow-calf enterprise — as long as reproductive performance of the heifers is not compromised.

Inconclusive results exist regarding the appropriate target weight, as well as the appropriate time heifers should be placed with bulls in relation to the mature cow herd. It is a common practice to breed heifers before the rest of the cow herd so they have a longer postpartum period before the start of their second breeding season. This management strategy requires additional resources if heifers are developed to the same target weight, because an accelerated rate of gain is needed to reach the target weight earlier.

There are studies comparing development of spring-born heifers at two prebreeding target weights (55% and 60% of mature weight) and the effects on reproduction and cow and calf productivity. Heifers on the high-gain development diet gained 0.3 pounds (lb.) per day more than heifers on the low-gain diet (1.4 vs. 1.1 lb. per day) and cost \$22 per head more during the 155-day wintering period.

At prebreeding, the high-gain heifers weighed 51 lb. more than the low-gain group and had 0.4-unit higher condition score. The high-gain heifers were 57% of mature weight and the low-gain heifers were 53% of mature weight at prebreeding. The percentage of heifers cycling before breeding was 11% higher for the high-gain heifers (85% vs. 74%). The 45-day pregnancy rate was not statistically different between groups (92% vs. 88% for low- and high-gain heifers, respectively).

► Developing heifers using crop residues may be a means to reduce development costs and not impair reproductive performance. Replacement heifers developed on crop residues will need to be supplemented to achieve targeted ADG.





Fig. 1: ADG of calves supplemented DDGS while grazing corn residue



Data for calving and weaning of the first calf, and reproduction (through fourth pregnancy), indicated high-gain heifers as cows were heavier at calving and weaning. These differences in heifer development weights carried over through the fourth pregnancy diagnosis.

Average calf birth date, calf birth weight, calving difficulty and calf losses were similar for both groups through three calf crops. Calf gain and adjusted 205-day weights were also similar for both groups of cows, indicating milk production was probably similar. Percentage of cows rebreeding for their second, third and fourth calves were similar for both groups. There were no differences in pregnancy or calf production data when heifers were developed at either 53% or 57% of mature weight. However, costs were increased for the high-gain group.

Using crop residues

Corn residues are a relatively inexpensive feed resource, but are low in protein and energy, especially for growing calves, backgrounded for entry into the feedlot or for summer pasture, or for replacement heifers. Beef producers often target a specific average daily gain (ADG), so it is important to know the amount of a supplement to feed to calves grazing corn residues in order to achieve a desired level of daily gain.

Due to their high energy [112% total digestible nutrients (TDN)] and high protein content (30.1%), dried distillers' grains with solubles (DDGS) fit well as a protein and/or energy supplement in many grazing situations. DDGS were fed to weanling steer calves grazing nonirrigated corn residue to determine daily gain response and residue intake response to increasing levels of DDGS (from 1.5 lb. to 6.5 lb. per day in 1-lb. increments; Gustad et al., 2006 *Nebraska Beef Report*; see Fig. 1). The DDGS was fed individually using Calan electronic gates. Daily gain increased from 0.9 lb. (1.5 lb. DDGS) per day to 1.8 lb. (6.5 lb. DDGS) per day. Forage intake decreased from 11.3 lb. (1.5 lb. DDGS) per day to 8.3 lb. (6.5 lb. DDGS) per day.

These results provide information for selecting a DDGS supplementation level to achieve a target gain. The ADGs in this experiment are similar to ADGs that many would target replacement heifers.

Final thoughts

These data would suggest that heifer development costs can be reduced by designing feeding programs to target heifers to 55%-60% of their mature weight before the start of the first breeding season. Heifers developed to lighter prebreeding weights maintained adequate breeding performance

during the second, third and fourth breeding seasons. If you consider this type of heifer development program, you may want to retain more heifers into the program in case low rebreeding performance is incurred and overall cow numbers can be maintained.

In addition, recently, females that do not get pregnant in their first breeding season have not been costly, as the price of yearlings sold after the grazing season has been good.

Finally, developing heifers using crop residues may be a means to reduce development costs and not impair reproductive performance. Replacement

heifers developed on crop residues will need to be supplemented to achieve targeted ADG as residues are low in protein and energy.

These management strategies can affect profit potential of the cow-calf enterprise.



E-MAIL: rrasby@unlnotes.unl.edu

Editor's Note: "Ridin' Herd" is a monthly column written by Rick Rasby, professor of animal science at the University of Nebraska. The column focuses on beef nutrition and its effects on performance and profitability.